

Polysaccharide Krestin Activity of Coriolus versicolor Extract on Interleukin-12 Level of Mus muculus Exposed to Mycobacterium tuberculosis

by Sri Puji Astuti Wahyuningsih

Submission date: 16-Nov-2018 04:07 PM (UTC+0800)

Submission ID: 1040211054

File name: Bukti_C18.pdf (422.23K)

Word count: 3637

Character count: 18888

1 **Polysaccharide Krestin Activity of *Coriolus versicolor***
2 **Extract on Interleukin-12 Level of *Mus musculus* Exposed**
3 **to *Mycobacterium tuberculosis***

4
5 **Sri Puji Astuti Wahyuningsih¹, Sugiharto¹, Nurul Wiqoyah² & Zakiyatun Nuha¹**

6 ¹Depart. of Biology, Faculty of Science and Technology, Airlangga University,
7 Surabaya 60115 Indonesia

8 ² Microbiology Laboratoty, Faculty of Medicine, Airlangga University, Surabaya 60115
9 Indonesia

10 Email: sri-p-a-w@fst.unair.ac.id sri-p-a-w@fst.unair.ac.id
11
12

13 **Abstract.** This study aimed to determine the effect of polysaccharide krestin (PSK) of
14 *Coriolus versicolor*'s extract on IL-12 levels in the *Mus musculus*'s blood serum that
15 had been exposed by *Mycobacterium tuberculosis*. This study used a completely
16 randomized design were divided into six treatment groups. They were K (control group
17 without treatment), K+ (positive control by providing PSK), K- (negative control with
18 exposed *M. tuberculosis*), P1 (treatment using PSK before exposed by *M.*
19 *tuberculosis*, P2 (treatment using PSK after exposed by *M. tuberculosis*), and P3
20 (treatment using PSK before and after exposed by *M. tuberculosis*). PSK given
21 concentration was 200 mg/kg bw, while the number of bacteria for exposure was 0.25
22 Mc Farland with double exposure. Each treatment there were four replicates. Blood
23 serum of mice were isolated and measured levels of IL-12 by ELISA kit. Data analysis
24 used Brown Forsythe test. The results showed that the highest levels of IL-4 was K-.
25 PSK administration in the treatment of P1, P2, and P3 showed the levels of IL-12 not
26 significant with the K and K+. Conclusion of the study was the treatment using PSK
27 had no effect on the level of IL-12 in *Mus musculus* exposed by *M. tuberculosis*.
28

29 **Keywords:** *Coriolus versicolor*; *interleukin-12*, *polysaccharides krestin*;
30 *Mycobacterium tuberculosis*.

31 **1 Introduction**

32 Tuberculosis or TB is an infectious bacterial disease caused by *Mycobacterium*
33 *tuberculosis*, which most commonly affects the lungs. It is transmitted from
34 person to person via droplets from the throat and lungs of people with the active
35 respiratory disease. The symptoms of active TB of the lung are coughing,
36 sometimes with sputum or blood, chest pains, weakness, weight loss, fever and
37 night sweats [Departemen Kesehatan RI, 7].
38

5
39 In healthy people, infection with *M. tuberculosis* Often causes no symptoms,
40 since the person's immune system acts to "wall off" the bacteria. Tuberculosis
41 can be treated with antibiotics for 6 months in a row. It is important to look for
42 other alternative materials that can be used to enhance the immune response to
43 tuberculosis.

3
44 Medicinal mushrooms have an established history of use in traditional oriental
45 therapies. Modern clinical practice in Japan, China, Korea, and other Asian
46 countries continues to rely on mushroom. Mushrooms effects have been
47 demonstrated for many including extracts of species from *C. versicolor* [Ooi &
48 Liu, 14].

1
49 It is well established that many mushroom-extracted compounds are commonly
50 used as immunomodulators or as Biological Response Modifiers (BRM). The
51 basic strategy underlying immunomodulation is to identify aspects of the host
52 response that can be enhanced or suppressed in such a way as to augment or
53 complement a desired immune response. Whether certain compounds enhance
54 or suppress immune responses depends on a number of factors, including dose,
55 route of administration, timing of administration of the compound, mechanism
56 of action, and site of activity. Knowledge of the specific components of
57 cytokine networks and signaling pathways and their role in the regulation of
58 immune responses is important in designing strategies to augment these
59 responses.

60 *C. versicolor* extract can increase the number of leukocytes, macrophages and
61 spleen weight [Wahyuningsih, 20], the provision of PSK from *C. versicolor*
62 increase the number of immunocompetent cells, increase immune response non-
63 specific and specific due to infection with *M. tuberculosis* (Wahyuningsih *et al.*,
64 21). The active compounds contained in mushrooms is β -glucan [Guggenheim,
65 9; Moradali *et al.*, 13].

66 β -Glucan is known to stimulate the formation of pro-inflammatory mediators
67 such as complement components, interleukin 1 (IL-1), tumor necrosis factor
68 (TNF- α), interleukin 2 (IL-2) and eicosanoids [Yu *et al.*, 27]. β -Glucan
69 increases the production of IL-2, which stimulates the differentiation of B cells
70 activated [Vetvicka *et al.*, 24]. In this study, administration of PSK done in
71 three different time ie before exposure, after exposure and before and after
72 exposure to *M. tuberculosis*. Giving PSK before exposure to *M. tuberculosis*
73 can serve as a preventative that will encourage the formation of antibodies so
74 the body does not easily infected. Giving PSK after exposure was conducted to
75 determine the effectiveness of the PSK in eliminating acid-resistant bacteria.
76 Giving PSK before and after exposure to *M. tuberculosis* as a preventive and
77 curative efforts that encourage the formation of antibodies [Wahyuningsih *et al.*,

Polysaccharide Krestin Activity of *C. versicolor* Extract on
Interleukin-12 Level of *M. mucus* Exposed to *M. tuberculosis* 3

78 22]. Pietro [15], β -glucan is effective for the prevention and treatment of
79 diseases related to the resilience of the body's immune system.

80 B-glucan compounds of PSK related to the primary receptor of the immune
81 system that is Dectin-1, Toll-Like Receptor 2/6 (TLR-2/6) and Complement
82 Receptor 3 (CR3). Dectin-1 known to play a role as a co-stimulator of T cells
83 and is the receptor that is expressed by macrophages [Willment, 23]. In addition
84 Dectin-1 can work with TLRs to induce and increase the production of
85 cytokines [Gantner *et al.*, 8]. Interleukin-12 binding to its receptor activates the
86 tyrosine kinase 2 (Tyk2) and janus kinase 2 (JAK2). This causes the
87 phosphorylation of tyrosine residues of the signal transducer and activator of
88 transcription 3 and 4 (STAT3 and STAT4). The occurrence of tyrosine
89 phosphorylation process is responsible for the formation of STAT4 / STAT4
90 homodimer and STAT3 / STAT4 heterodimers. Both dimers are experiencing
91 translocation into the nucleus and binds to the IFN- γ gene promoter [Thierfelder
92 *et al.*, 19]. Promoter activation in the IFN- γ gene resulted in the synthesis of
93 proteins that express IFN- γ protein.

94 Along with the growing understanding of the body's immune response in the
95 face of infection, the more developed also research into components that can
96 affect the immune response. If PSK can increase the levels of IL-12 in the
97 blood, the immune system can increase too. If it is proven that extracts of *C.*
98 *versicolor* can be used as a natural immunostimulatory to inhibit the growth of
99 *M. tuberculosis* and reduce the number of TB patients. Active TB disease is
100 estimated to have reached 1.9 billion, and it can prevent transmission because
101 only people with active TB can transmit the disease.

102 2 Materials and Methods

103 2.1 Preparation research

104 Thirty female mice aged 8-10 weeks strain Balb/C acclimatized for one week
105 and are grouped into six, namely K (only given distilled water), K- (exposure to
106 *M. tuberculosis* only), K + (given PSK only), P1 (PSK was given before
107 exposure to *M. tuberculosis*), P2 (PSK administered after exposure to *M.*
108 *tuberculosis*), P3 (PSK given before and after exposure to *M. tuberculosis*).
109 Feed and water provided ad libitum.

110 2.2 Isolation and Measurement Levels PSK

111 *Coriolus versicolor* were collected from Kediri, Tulungagung and Surabaya.
112 Fungi identified, wind dried and pulverized to form a coarse powder. Then it

113 was extracted by the method of Cui & Cristi [5] and Cui *et al.* [6] as modified
114 by Wahyuningsih *et al.* [20] PSK levels determined by the method of phenol-
115 sulfuric acid assay. PSK dose used was 200 mg/Kg body weight.

116 2.3 Treatment research

117 Giving PSK done for 7 days. Exposure to *M. tuberculosis* performed twice with
118 an interval of two weeks via intraperitoneal. The number of bacteria is 0.5 Mc
119 Farland.

121 2.4 Isolation serum

122 Blood was drawn through intracardia. Blood left in a tilted position at room
123 temperature for three hours to form two phases, namely the colored translucent
124 top and the bottom is red. Then, it centrifuged 3000 rpm, 10 min at 4 ° C. Serum
125 or supernatant portion was taken.

126 2.5 Immunosorbent-Like Enzyme Assay (ELISA)

127 Detection of IL-12 used to a sandwich ELISA (Koma Biotech ELISA Kit IL-
128 12). Levels of IL-12 was read by ELISA reader at a wavelength of 450 nm.

129 2.6 Data analysis

130 Data were tested with the Kolmogorov-Smirnov test and homogeneity of
131 variance. Then analyzed by Brown Forsythe at the level of 5% ($\alpha = 0.05$).

132 3 Result

133 Levels of Interleukin 12 in the blood serum is a picture of an immune response
134 as a result of exposure to *M. tuberculosis* were shown in Table 1. The results of
135 the ELISA is a yellow color reaction. This color is obtained after a complete
136 bond between the antigen, primary antibody and enzyme-labeled secondary
137 antibody.

138 The results showed that average levels of IL-12 in the control group (K) was
139 25.91 ± 11.08 (pg / mL). Positive control group (K +) was 26.37 ± 11.49 (pg /
140 mL). Negative control group (K) was 37.25 ± 27.73 (pg / mL). P1 group was
141 31.85 ± 16.22 (pg / mL). P2 group was 31.43 ± 10.47 (pg / mL). P3 group was
142 26.07 ± 10.99 (pg / mL). Kolmogorov-Smirnov test showed a significance level
143 of 0.068. This indicates that the data were normally distributed. Homogeneity
144 test showed a significance level of 0,010. This indicates that the data has a

Polysaccharide Krestin Activity of *C. versicolor* Extract on Interleukin-12 Level of *M. mucus* Exposed to *M. tuberculosis* 5

variance that is not homogeneous. Data were analyzed by Brown Forsythe test to determine the effect of treatment in each group. The test results showed a significance level of 0.828. This shows that there was no effect of feeding time on levels of IL-12 in mice blood serum as a result of exposure to *M. tuberculosis*.

Table 1 Levels of IL-12 after administration of PSK from *C. versicolor* in mice exposed to *M. tuberculosis*

Treatment	Levels of IFN- γ on replay to				Mean \pm SD (pg/mL)
	1	2	3	4	
K	24.90	30.75	27.32	20.68	25.91 ^a \pm 11.08
K+	30.44	31.07	23.28	20.68	26.37 ^a \pm 11.49
K-	81.88	26.90	19.84	20.36	37.25 ^a \pm 27.73
P1	33.57	48.90	23.04	21.89	31.85 ^a \pm 16.22
P2	24.77	32.05	45.97	22.93	31.43 ^a \pm 10.47
P3	21.22	26.90	29.66	26.48	26.07 ^a \pm 10.99

4 Discussion

In the study used an indicator to determine the levels of cytokine immune response to infection with *M. tuberculosis* and also the effect of PSK. Cytokines are small proteins as a mediator and regulator of immunity, inflammation and hematopoiesis. The cytokines produced in response to the stimulus of the immune system. Cytokines works by binding to specific membrane receptors, which then carry signals into the cell via second messengers (tyrosine kinase), to alter gene expression (Judarwanto, 11). Cytokines produced in the body when the immune response as a result of infectious microorganisms or foreign substances that enter the body (Romagnani, 16). Type of cytokine that is used as an indicator in this study is the production of Th1 cytokines, namely IL-12. Interleukin is a cytokine that acts against leukocytes, while interferon is a type of cytokine that plays a role due to interference virus (Lewis, 12).

Tuberculosis is a disease caused by intracellular bacterial pathogens, namely *M. tuberculosis* (Higuchi et al., 10). One type of cytokine that plays an important role in the defense against intracellular pathogens such as *M. tuberculosis* is IL-12. APC cells produce IL-12 when stimulated by lypotechoat acid and peptidoglycan. Two components are bacterial cell wall constituent. Wall of *M. tuberculosis* can induce macrophages to produce IL-12 (Yoshida & Koide, 26).

174 In this study, *M. tuberculosis* acts as an antigen. The bacteria will induce the
175 secretion of IL-12. PSK act as immunostimulatory which is expected to enhance
176 the immune response due to infection with *M. tuberculosis*. Polysaccharides
177 krestin from *C. versicolor* extract known to contain β -glucan compounds. The
178 compounds may be associated with the primary receptor of the immune system
179 such as Dectin-1, Toll-Like Receptor 2/6 (TLR-2/6) and Complement Receptor
180 3 (CR3). Dectin-1 acts as a co-stimulator of T cells and is the receptor that is
181 expressed by macrophages (Willment, 23).

182 In Table 1 it can be seen that the levels of IL-12 between the control and
183 treatment groups showed no significant difference. However, if compared to K,
184 then the levels of IL-12 on K +, K-, P1, P2, and P3 showed an increase. Levels
185 of IL-12 highest detected in the K group treated only exposure to *M.*
186 *tuberculosis*. It shows that there are components of antigens on the cell wall of
187 bacteria that can enhance the immune response. According to Crick *et al.* (4),
188 TB bacterial cell wall composed of mannose-capped lipoarabinomannan
189 (ManLAM), lipomannan (LM), phosphatidyl-myo-inositol mannosides (PIMs),
190 arabinomannan, mannan and Manno-glycoproteins. It is a strong antigen to
191 induce an immune response. Through TLRs, *M. tuberculosis* cell wall
192 components associated with lipoprotein so as to induce the production of IL-12
193 (Brightbill *et al.*, 2).

194
195 PSK was known to induce the body's immune response, but the strength of
196 induction was weaker when compared to components of *M. tuberculosis*
197 antigens. It can be seen from Table 1, which showed that levels of IL-12 to K +
198 is higher when compared with K. Levels of IL-12 in the group P1, P2, and P3
199 showed lower than K-. This was probably related to the role of β -glucan.
200 Glucan can bind directly to specific receptors of immune cells causing
201 immunomodulating effects (Vos *et al.*, 25). This means it can increase or
202 decrease the immune response. But the mechanism of action of β -glucan as
203 immunomodulators, especially in *M. tuberculosis* infection is not known.

204
205 Statistically, the provision PSK does not affect the levels of IL-12. Most likely
206 behind it was the existence of mechanisms mediated cascade of other cytokines
207 stimulated due to the antigen of *M. tuberculosis*. Another possibility was
208 happening related to the nature of cytokine that a type of cytokine can be
209 produced by various types of cells and can cause different effects through
210 various channels (Soeroso, 17). Various types of cytokine sometimes also have
211 the same function for one type of cells (Abbas *et al.*, 1). The redundancy
212 properties that make it difficult to know the mechanism of action of one single
213 type of cytokine.

Polysaccharide Krestin Activity of *C. versicolor* Extract on Interleukin-12 Level of *M. mucus* Exposed to *M. tuberculosis* 7

214 β -Glucan contained in PSK of *C. versicolor* known to play a role as an
215 immunomodulator that acts on the immune system of specific and non-specific.
216 In the non-specific immune system, β -glucan helps to recognize the antigen and
217 respond faster. Dectin-1 is a receptor type II transmembrane protein, can bind to
218 β -1,3 and β -1,6 glucan that can do the initiation and regulation of the non-
219 specific immune response (Brown *et al.*, 3). Dectin-1 is expressed by immune
220 cells that play a role in non-specific immune system and have been found on
221 macrophages, neutrophils, and dendritic cells (Taylor *et al.*, 18). In the specific
222 immune response, Dectin-1 is able to activate T cells and induces the secretion
223 of cytokines (Taylor *et al.*, 18). According to Gantner *et al.* (8), Dectin-1 is able
224 to induce and increase the secretion of cytokines through TLRs. In addition, the
225 β -glucan known to be effective for the prevention and treatment of diseases
226 associated with immune system resistance (Pietro, 15).

227 The conclusion from this study is the provision PSK dose of 50 mg/kg body
228 weight had no effect on levels of IL-12 in mice blood serum as a result of
229 exposure to *M. tuberculosis*. Further studies, using other indicators that can be
230 known immune mechanisms of TB patients by administration of PSK from *C.*
231 *versicolor* extracts.

232 5 References

- 233 [1] Abbas, A.K., Lichtman, A.H., & Pober, J.S., *Cytokine in cellular and*
234 *molecular immunology*, WB Saunders Co, Philadelphia, 1994.
- 235 [2] Brightbill, H.D., Libraty, D.H., Krutzik, S.R., Yang, R.B., Belisle, J.T.,
236 Bleharski, J.R., Maitland, M., Norgard, M.V., Plevy, S. E., Smale, S.T.,
237 Brennan, P. J., Bloom, B.R., Godowski, P.J., & Modlin, R.L., *Host*
238 *defense mechanisms triggered by microbial lipoproteins through toll-like*
239 *receptors*, Science, 285:732–736, 1999.
- 240 [3] Brown, G.D., Herre, J., Williams, D.L., Willment, J.A., Marshall, A.S., &
241 Gordon, S., *Dectin-1 mediates the biological effects of beta-glucans*, *The*
242 *journal of experimental medicine*, 197(9): 1119-1124, 2003.
- 243 [4] Crick, D.C., Brennan, P.J., & McNeil, M.R., *The cell wall of*
244 *Mycobacterium tuberculosis*, Lippincott Williams and Wilkins,
245 Philadelphia, 2003.
- 246 [5] Cui, J. & Chisti, Y., *Polysaccharopeptides of Coriolus versicolor:*
247 *physiological activity, uses, and production*, *Biotechnology advances*, 21
248 (2):109-122, 2003,.
- 249 [6] Cui, J., Goh, K.K.T., & Archer, R., *Characterisation and bioactivity of*
250 *protein-bound polysaccharides from submerged-culture fermentation of*
251 *Coriolus versicolor Wr-74 and ATCC-20545 strains*, *Journal Industrial*
252 *Microbiology Biotechnology*, 34: 393–402, 2007.

- 253 [7] Departemen Kesehatan Republik Indonesia, *TBC*, Pusat Komunikasi
254 Publik Setjen Depkes RI, Jakarta, 2008.
- 255 [8] Gantner, B.N., Simmons, R.M., Canavera, S.J., Akira, S., & Underhill,
256 D.M., *Collaborative induction of inflammatory responses by dectin-1 and*
257 *Toll-like receptor 2*, Journal of experimental medicine, 197:1107–1117,
258 2003.
- 259 [9] Guggenheim, A.G., Wright, K.M., & Zwickey, H.L., *Immune modulation*
260 *from five major mushrooms: Application to integrative oncology,*
261 *Integrative medicine*, 13: 32, 2014.
- 262 [10] Higuchi, K., Sekia, Y., & Harada, N., *Characterization of M.*
263 *Tuberculosis-derived IL-12-inducing material by alveolar macropages,*
264 *Vaccine*, 22: 724-734, 2003.
- 265 [11] Judarwanto, W., *Imunologi dasar: Sitokin dan aspek klinisnya,*
266 <http://allergycliniconline.com>, 2012.
- 267 [12] Lewis, R., *Cytokine*, <http://www.the-scientist.com> (14 Januari 2015).
- 268 [13] Moradali, M.F., Mostafavi, H., Ghods, S., & Hedjaroude, G.A.,
269 *Immunomodulating and anticancer agents in the realm of macromycetes*
270 *fungi (macrofungi)*. *Int immunopharmacology*. 7: 701-724, 2007,.
- 271 [14] Ooi, V.E & F. Liu, *Immunomodulation and Anticancer activity of*
272 *Polysaccharide - Protein Complexes*, National Library of Medicine,
273 *CurrMedChem*, <http://www.ncbi.nlm.nih.gov/pubmed/10702635> (4 Mei
274 2009)
- 275 [15] Pietro, P. , *Composition for preventif and/or treatment of lipid*
276 *metabolism disorders and allergic form*. <http://freepatentonline.com>. (20
277 April 2014).
- 278 [15] Romagnani, S., Lymphokine production by human T cells in disease
279 states. *Annu review immunology*, 12: 227-257, 1994.
- 280 [17] Soeroso, A., Sitokin, *Jurnal oftalmologi Indonesia*, 5: 171-180, 2007.
- 281 [18] Taylor, P.R., Brown, G.D, Harre, J., Williams, D.S., Willment, J.A., &
282 Gordon, S., *The role of SIGNR I and the beta-glucan receptor (dectin-I)*
283 *in the nonopsonic recognition of yeast by specific macropages*, *Journal*
284 *immunology*, 172(2): 1157-1162, 2004.
- 285 [19] Thierfelder WE, van Deursen JM, Yamamoto K, Tripp RA, Sarawar SR,
286 Carson RT, Sangster MY, Vignali DA, Doherty PC, Grosveld GC, & Ihle
287 JN, *Requirement for Stat4 in interleukin-12-mediated responses of*
288 *natural killer and T cells*, *Nature*, 382: 171, 1996.
- 289 [20] Wahyuningsih, S.P.A., *Pemanfaatan ekstrak jamur Coriolus versicolor*
290 *sebagai imunomodulator respon imun non-spesifik pada tikus putih*
291 *akibat infeksi M. tuberculosis*, Lembaga Penelitian Universitas Airlangga,
292 Penelitian DIPA Universitas Airlangga, Surabaya, 2006.
- 293 [21] Wahyuningsih, S.P.A., Darmanto, W., & Wiqoyah, N., *Bioaktivitas*
294 *polisakarida krestin dari ekstrak jamur Coriolus versicolor sebagai*
295 *imunomodulator respon imun akibat infeksi Myobacterium tuberculosis,*

Polysaccharide Krestin Activity of *C. versicolor* Extract on
Interleukin-12 Level of *M. mucus* Exposed to *M. tuberculosis* 9

- 296 Lembaga Penelitian Universitas Airlangga, Hibah Penelitian Strategis
297 Nasional, Surabaya, 2009.
- 298 [22] Wahyuningsih, S.P.A., Winarni, D., & Masitha, A.N., *Aktivitas*
299 *polisakarida krestin dari ekstrak Coriolus versicolor terhadap*
300 *peningkatan antibodi Mus musculus akibat paparan Mycobacterium*
301 *tuberculosis*, Berkala penelitian hayati, 17: 177-183, 2012.
- 302 [23] Willment, J.A., *The human beta-glucan receptor is widely expressed and*
303 *functionally equivalent to murine Dectin-1 on primary cells*, Eur journal
304 *immunology*, 5: 1539–1547, 2005.
- 305 [24] Vetvicka, V., Kiyomi, T., Rosemade, M., Paulin, B., Bill, K., & Gary, O.
306 *Orally-Administered Yeast β -1,3 glucan Prophylactically Protects*
307 *Against Anthrax Infection and Cancer in Mice*, Journal American
308 Nutraceutical Assosiation, 5: 2, , 2002.
- 309 [25] Vos, A., M'Rabet, L., Stahl, B., Boehm, G., & Garssen, J., *Immune-*
310 *modulatory effects and potential working mechanisms of orally applied*
311 *nondigestible carbohydrates*, Crit review immunology, 27: 97-140, 2007.
- 312 [26] Yoshida, A., & Koide, Y., *Arabino-furanosyl-terminated and*
313 *mannosylated lipoarabinomannans from Mycobacterium tuberculosis*
314 *induce different levels of interleukin-12 expression in mouse*
315 *macrophages*, Infect immun, 65: 1953–1955, 1997.
- 316 [27] Yu, S., Weaver, V., Martin, K., & Cantorna, M.T., *The effects of whole*
317 *mushrooms during inflammation*, BMC Immunology, 10:12, 2009.

Polysaccharide Krestin Activity of Coriolus versicolor Extract on Interleukin-12 Level of Mus musculus Exposed to Mycobacterium tuberculosis

ORIGINALITY REPORT

16%

SIMILARITY INDEX

9%

INTERNET SOURCES

12%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

1

Ben-Zion Zaidman. "Medicinal mushroom modulators of molecular targets as cancer therapeutics", Applied Microbiology and Biotechnology, 06/2005

Publication

4%

2

www.countyofsb.org

Internet Source

2%

3

www.coriolusversicolor.us

Internet Source

1%

4

lsresearch.thomsonreuters.com

Internet Source

1%

5

globalhealthtrainingcentre.tghn.org

Internet Source

1%

6

"Researchers Submit Patent Application, "Combination Therapy", for Approval.", Biotech Week, April 8 2015 Issue

Publication

1%

7	Torrelles, J.B.. "Diversity in Mycobacterium tuberculosis mannosylated cell wall determinants impacts adaptation to the host", Tuberculosis, 201003 Publication	1 %
8	healty7611.blogspot.my Internet Source	<1 %
9	www-ermm.cbcu.cam.ac.uk Internet Source	<1 %
10	www.freepatentsonline.com Internet Source	<1 %
11	Polysaccharides, 2015. Publication	<1 %
12	Olsson, S.. "The macrophage @b-glucan receptor mediates arachidonate release induced by zymosan: Essential role for Src family kinases", Molecular Immunology, 200703 Publication	<1 %
13	lib.ugent.be Internet Source	<1 %
14	Venema, Koen. "In Vitro Assessment of the Bioactivity of Food Oligosaccharides", Food Oligosaccharides, 2014. Publication	<1 %

15

epdf.tips
Internet Source

<1 %

16

Finimundy, Tiane Cristine, Aldo José Pinheiro Dillon, João Antônio Pêgas Henriques, and Mariana Roesch Ely. "A Review on General Nutritional Compounds and Pharmacological Properties of the *Lentinula edodes* Mushroom", Food and Nutrition Sciences, 2014.

Publication

<1 %

17

orbi.ulg.ac.be
Internet Source

<1 %

18

"Evidence-based Anticancer Materia Medica", Springer Nature America, Inc, 2011

Publication

<1 %

19

ajprenal.physiology.org
Internet Source

<1 %

20

mobile.repository.ubn.ru.nl
Internet Source

<1 %

21

Dietary Components and Immune Function, 2010.

Publication

<1 %

22

Kathrin Luther. "Phagocytosis of *Aspergillus fumigatus* conidia by murine macrophages involves recognition by the dectin-1 beta-glucan receptor and Toll-like receptor 2",

<1 %

23

Sun, Lina, and Yong Zhao. "The Biological Role of Dectin-1 in Immune Response", International Reviews of Immunology, 2007.

Publication

<1%

24

Principles of Immunopharmacology, 2011.

Publication

<1%

25

"Hepatitis C Virus Disease", Springer Nature America, Inc, 2008

Publication

<1%

Exclude quotes

Off

Exclude matches

Off

Exclude bibliography

On

Polysaccharide Krestin Activity of Coriolus versicolor Extract on Interleukin-12 Level of Mus mucus Exposed to Mycobacterium tuberculosis

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9